arXiv:0711.3385v1 [math.DS] 21 Nov 2007

Global attractor in competitive Lotka-Volterra systems

Zhanyuan Hou

Abstract

For autonomous Lotka-Volterra systems of differential equations modelling the dynamics of n competing species, new criteria are established for the existence of a single point global attractor. Under the conditions of these criteria, some of the species will survive and stabilise at a steady state whereas the others, if any, will die out.

Keywords Lotka-Volterra, global attractor, competitive systems
MSC(2000) 34D45, 92D25

1 Introduction

In this paper, we are concerned with the asymptotic behaviour of solutions of the autonomous Lotka-Volterra system

$$x_i' = b_i x_i(1 - \alpha_i x), \quad i \in I_n, \tag{1.1}$$

where $x_i' = dx_i/dt$, $b_i > 0$, $I_m = \{1, 2, \ldots, m\}$ for any positive integer m, $\alpha_i = (a_{i1}, a_{i2}, \ldots, a_{in})$ with $a_{ii} > 0$ and $a_{ij} \geq 0$, and $x = (x_1, x_2, \ldots, x_n)^T \in \mathbb{R}^n$. Since the behaviour of (1.1) is clear when $n = 2$, we assume $n > 2$ though some of our results given later are applicable to two-dimensional systems. Let $\mathbb{R}^n_+ = \{x \in \mathbb{R}^n : x \geq 0\}$, where by $y \geq x$ in $\mathbb{R}^n$ we mean $y_i \geq x_i$ for all $i \in I_n$. We also write $y > x$ if $y \geq x$ but $y \neq x$. We shall restrict the solutions of (1.1) to $\mathbb{R}^n_+$ in view of (1.1) modelling the dynamics of n competitive species, $x_i(t)$ denoting the population size of the ith species at time t. We say that an equilibrium $x^* \in \mathbb{R}^n_+$ of (1.1) is a global attractor if every solution of (1.1) in $\text{int}\mathbb{R}^n_+ = \{x \in \mathbb{R}^n : x_i > 0, i \in I_n\}$, i.e. with $x(0) \in \text{int}\mathbb{R}^n_+$, satisfies $\lim_{t\to\infty} x(t) = x^*$.

Since the independent introduction of (1.1) as a population model by Lotka and Volterra in the 1920s, a large number of researchers have been attracted to the theoretical investigation of (1.1) and its variations (see, for example, [13] and the references therein for a brief account of the development). One of the problems arising from (1.1) as a model is to find suitable conditions under which some particular (or all) species in the competitive community will survive and stabilise at a steady state whereas the rest, if any, will die out eventually. Mathematically, we need suitable conditions which ensure the existence of an equilibrium $x^* \in \mathbb{R}^n_+$ that is a global attractor. It is known that (1.1) has an equilibrium $x^* \in \text{int}\mathbb{R}^n_+$ that is a global attractor if

$$\frac{a_{i1}}{a_{11}} + \frac{a_{i2}}{a_{22}} + \cdots + \frac{a_{in}}{a_{nn}} < 2 \tag{1.2}$$

holds for all $i \in I_n$ (see [10], [6] or [5, p. 294–297]). For (1.1) to have the same property as above, Hou [7] finds a more general condition

$$\max\left\{0, \frac{a_{ij}}{a_{jj}}\left(1 - \sum_{k\neq i,j} \frac{a_{jk}}{a_{kk}}\right)\right\} < 1 - \sum_{k\neq i,j} \frac{a_{ik}}{a_{kk}} \tag{1.3}$$

for all $i, j \in I_n$ with $i \neq j$, which incorporates (1.2) for all $i \in I_n$ as a special case. For (1.1) to have an equilibrium $x^* \in \partial\mathbb{R}^n_+ = \mathbb{R}^n_+ \setminus \text{int}\mathbb{R}^n_+$ that is a global attractor, Zeeman [14] shows that

$$(i-j)(a_{ij}-a_{jj}) > 0 \tag{1.4}$$

for $i, j \in I_n$ with $i \neq j$ guarantees the survival of only the first species, and Ahmad and Lazer [1] obtain a criterion for the extinction of only one species. Li, Yu and Zeng [11] extend this criterion to the extinction of possibly more than one species. Combining (1.3) with a condition similar to that of [12] for nonautonomous systems, Hou [7] shows the global attraction of an equilibrium $x^* \in \partial\mathbb{R}^n_+$ with the survival of some species and the extinction of the others.

The main purpose of this paper is to establish new criteria for the existence of a single point global attractor. Using the idea of the "ultimate contracting cells" method, which was summarised by Hou [8] or [9], rather than using Lyapunov functions, we shall see that the new conditions are geometrically simple and include (1.2), (1.3) and those given in [1] as particular instances.

The presentation of this paper is as follows. The statements of the main results under geometric conditions, together with some examples, will be given in §2. Then in §3 these geometric conditions will be translated into algebraic expressions. In §4 the main results will be restated under equivalent algebraic conditions. Finally, the proofs of the main theorems will be completed in §5.

2 Main results under geometric conditions

We first explain some terms and notation. For any $u, v \in \mathbb{R}^n_+$ with $v \geq u$, a *cell* $[u, v]$ is defined to be the set of $x \in \mathbb{R}^n_+$ satisfying $v \geq x \geq u$. For any $J \subset I_n$, $u^J \in \mathbb{R}^n_+$ is defined by $u^J_i = u_i$ if $i \in J$ and $u^J_i = 0$ otherwise. Thus, with this notation, $u^{I_n} = u$, $u^\emptyset = 0$, and $u^{I_n \setminus \{k\}}$ for any $k \in I_n$ is the projection of u onto the coordinate plane π_k, which is defined by (2.2). If $u > 0$, the set $\gamma = \{x \in \mathbb{R}^n_+ : u^T x = 1\}$ is an $(n-1)$-dimensional *plane* in $\mathbb{R}^n_+$. A point $x \in \mathbb{R}^n_+$ is said to be *below* (*on* or *above*) γ if $u^T x < 1$ ($u^T x = 1$ or $u^T x > 1$). A nonempty set $S \subset \mathbb{R}^n_+$ is said to be *below* (*on* or *above*) γ if every point x in S is so. Now for (1.1) we let

$$Y = (a_{11}^{-1}, a_{22}^{-1}, \ldots, a_{nn}^{-1})^T, \tag{2.1}$$

$$\pi_i = \{x \in \mathbb{R}^n_+ : x_i = 0\}, \quad i \in I_n, \tag{2.2}$$

$$\gamma_i = \{x \in \mathbb{R}^n_+ : \alpha_i x = 1\}, \quad i \in I_n, \tag{2.3}$$

and view $\cap_{i\in\emptyset}\pi_i$ as $\mathbb{R}^n_+$. Then, for any $J \subset I_n$, $\cap_{i\in J}\pi_i$ is obviously invariant under (1.1). Thus, as $\gamma_i \cap (\cap_{j\in I_n\setminus\{i\}}\pi_j) = \{Y^{\{i\}}\}$, $Y^{\{i\}}$ is an equilibrium of (1.1) on the x_i-axis. Let $U \in [0, Y]$ be defined as follows: for each $i \in I_n$, if γ_i is below γ_j for every $j \in I_n \setminus \{i\}$ then $U_i = 0$; if $Y^{\{i\}}$ is on or above some γ_j ($j \neq i$) then $U_i = Y_i = a_{ii}^{-1}$; otherwise U_i is the maximum value of the ith coordinate components of the intersection points of γ_i with any other γ_j.

CONDITION (C_k)**.** Either $U^{I_n\setminus\{k\}}$ is below γ_k or $\gamma_k \cap [0, U^{I_n\setminus\{k\}}]$ is above γ_j for every $j \in I_n \setminus \{k\}$. Alternatively, either $U^{I_n\setminus\{k\}}$ is below γ_k or the set $\gamma_j \cap [0, U^{I_n\setminus\{k\}}]$ is below γ_k for every $j \in I_n \setminus \{k\}$.

The analytical definition of U, as well as its simplification, and algebraic realisation of condition (C_k) are left to the next section. We shall see in §5 that U is an ultimate upper




Figure 2.1. $U^{I_3\setminus\{i\}}$ is below γ_i for $i \in I_3$, but $\gamma_3 \cap [0, Y^{\{2,3\}}]$ does not below γ_1.

bound for every solution $x(t)$ of (1.1) in $\mathrm{int}\mathbb{R}^n_+$, i.e. $\overline{\lim}_{t\to\infty} x_i(t) \le U_i$ for all $i \in I_n$. Moreover, condition (C_k) will guarantee the existence of some $\delta > 0$ such that every solution of (1.1) in $\mathrm{int}\mathbb{R}^n_+$ satisfies $\underline{\lim}_{t\to\infty} x_k(t) \ge \delta$. Most importantly, the following theorem holds.

Theorem 2.1 *Assume that (C_k) holds for all $k \in I_n$. Then (1.1) has a unique equilibrium $x^* \in \mathrm{int}\mathbb{R}^n_+$ that is a global attractor. Moreover, the inequality $x^* \le U$ holds.*

Remark 2.2 It is shown in [7] that (1.2) for all $i \in I_n$ implies (1.3) for all $i, j \in I_n$ with $i \ne j$. It is also shown that (1.3) for all $i, j \in I_n$ with $i \ne j$ is equivalent to that $\gamma_j \cap [0, Y^{I_n\setminus\{k\}}]$ is below γ_k for all $j, k \in I_n$ with $j \ne k$. Since $U \le Y$, (1.3) for all $i, j \in I_n$ with $i \ne j$ implies condition (C_k) for all $k \in I_n$. Therefore, theorem 2.1 incorporates the known results using (1.2) and (1.3) as special instances. The example below satisfies (C_k) for all $k \in I_n$ but does not satisfy (1.3) for some $i, j \in I_n$. Hence, theorem 2.1 applies to a broader class of systems than the known results do.

Example 2.3 Consider system (1.1) with $n = 3$ and

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} = \begin{pmatrix} 2.1 & 1.9 & 1.9 \\ 1 & 3 & 1 \\ 1 & 1 & 3 \end{pmatrix}. \tag{2.4}$$

Then $a_{ii} > a_{ji}$ holds for $i, j \in I_3$ with $i \ne j$ so $Y^{\{i\}}$ is below γ_j for all $i, j \in I_3$ with $i \ne j$. Since $\gamma_1 \cap \gamma_2 \cap \pi_3 = \{(0.25, 0.25, 0)^T\}$, $\gamma_1 \cap \pi_2 \cap \gamma_3 = \{(0.25, 0, 0.25)^T\}$, $\pi_1 \cap \gamma_2 \cap \gamma_3 = \{(0, 0.25, 0.25)^T\}$ and $(\cap_{j\in I_3\setminus\{i\}} \gamma_j) \cap \pi_i$ is below γ_i for $i \in I_3$, by definition we have $U = (0.25, 0.25, 0.25)^T$. Note that $\alpha_i U^{I_3\setminus\{i\}} < 1$, so (C_i) holds, for all $i \in I_3$ (see Figure 2.1). By theorem 2.1, $\gamma_1 \cap \gamma_2 \cap \gamma_3 = \{x^*\}$ with $U_i \ge x^*_i > 0$ for $i \in I_3$ and x^* is a global attractor. Indeed, $x^* = (\frac{1}{23}, \frac{11}{46}, \frac{11}{46})^T$. On the other hand, however, for $Y = (\frac{10}{21}, \frac{1}{3}, \frac{1}{3})^T$, we have $(0, \frac{1}{3}, \frac{11}{57})^T \in \gamma_1 \cap [0, Y^{\{2,3\}}]$ but $(0, \frac{1}{3}, \frac{11}{57})^T$ is below γ_3. So (1.3) does not hold for some $i, j \in I_3$.

Corollary 2.4 *Assume that (C_k) holds for all $k \in I_n$. Then, for each nonempty set $J \subset I_n$, $(\cap_{j\in J} \gamma_j) \cap (\cap_{\ell\in I_n\setminus J} \pi_\ell) = \{u_J\}$ with $U_j \ge (u_J)_j > 0$ for $j \in J$ such that every solution of (1.1) with $x_j(0) > 0$ for $j \in J$ and $x_\ell(0) = 0$ for $\ell \in I_n \setminus J$ satisfies $\lim_{t\to\infty} x(t) = u_J$. Moreover, (1.1) has 2^n equilibria in $\mathbb{R}^n_+$.*

Proof. When $J = \{i\}$ for some $i \in I_n$, (1.1) on $\cap_{k\in I_n\setminus\{i\}} \pi_k$ is equivalent to $x_i' = b_i x_i(1 - a_{ii}x_i)$ so $u_J = Y^{\{i\}}$ meets the requirement. Note that theorem 2.1 can be applied to (1.1) with $n > 1$.

Then, if $|J| > 1$, the conclusion about the required u_J follows from the application of theorem 2.1 to the $|J|$-dimensional system (1.1) on $\cap_{\ell\in I_n\setminus J}\pi_\ell$. Clearly, u_J is the unique equilibrium of (1.1) in $\cap_{\ell\in I_n\setminus J}\pi_\ell$ satisfying $(u_J)_j > 0$ for $j \in J$. Since I_n has $2^n - 1$ nonempty subsets and the origin is also an equilibrium, (1.1) has 2^n equilibria in $\mathbb{R}^n_+$.

We have mentioned earlier that condition (C_k), for any particular $k \in I_n$, will secure the survival of the kth species. Bearing this in mind and applying condition (C_k) to some of the indices (not necessarily all), we have the following.

Theorem 2.5 *Assume that condition (C_k) holds for all $k \in J \subset I_n$ $(J \neq \emptyset)$. Then $(\cap_{j\in J}\gamma_j) \cap (\cap_{k\in I_n\setminus J}\pi_k) = \{x^*\}$ with $0 < x^*_j \leq U_j$ for $j \in J$. In addition, if x^* is on γ_k for every $k \in I_n \setminus J$, then x^* is a global attractor. Moreover, the conclusion still holds if "x^* is on γ_k" is replaced by "x^* is on or above γ_k" when U is replaced by Y.*

Remark 2.6 Theorem 2.1 is the extreme case of theorem 2.5 when $J = I_n$. When $J = I_{n-1}$, Ahmad and Lazer [1] obtained the same conclusion under the assumptions that (i) (1.2) holds for all $i \in I_{n-1}$, (ii) $\alpha_n x^* > 1$ and (iii) $\det(a_{ij})_{n\times n} > 0$. Later in [2, 3] it was pointed out that the above condition (iii) is redundant. From remark 2.2 we know that, for any $i \in I_n$, (1.2) implies (1.3) for all $j \neq i$, which ensures condition (C_i) with the replacement of U by Y. Thus, the result in [1] under the assumptions (i) and (ii) is covered by theorem 2.5 as a special case.

Example 2.7 Consider system (1.1) with $n = 4$ and

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{pmatrix} = \begin{pmatrix} 5/2 & 3/2 & 1 & 3/2 \\ 1 & 3 & 1 & 1 \\ 2 & 2 & 2 & 2 \\ 2 & 2 & 1 & 3 \end{pmatrix}.$$

Then $Y = (\frac{2}{5}, \frac{1}{3}, \frac{1}{2}, \frac{1}{3})^T$. Clearly, $Y^{\{i\}}$ is below γ_j for every $j \in I_4 \setminus \{i\}$ as $a_{ii} > a_{ji}$. To calculate U_1, we find all the possible intersection points of γ_1 with γ_j $(j \neq 1)$ on $\pi_2 \cap \pi_3$, $\pi_2 \cap \pi_4$ and $\pi_3 \cap \pi_4$ and find the maximum of the x_1 values of these points. On $\pi_3 \cap \pi_4$, γ_1 intersects γ_j at $(\frac{1}{4}, \frac{1}{4}, 0, 0)^T$ for all $j \in I_4 \setminus \{1\}$. On $\pi_2 \cap \pi_4$, γ_1 intersects γ_3 at $(\frac{1}{3}, 0, \frac{1}{6}, 0)^T$; γ_1 intersects γ_2 and γ_4 on π_1 so these are ignored. On $\pi_2 \cap \pi_3$, γ_1 intersects γ_3 at $(\frac{1}{4}, 0, 0, \frac{1}{4})^T$ and γ_4 at $(\frac{1}{3}, 0, 0, \frac{1}{9})^T$; γ_1 is below γ_2. Then $U_1 = \max\{\frac{1}{4}, \frac{1}{3}\} = \frac{1}{3}$. We then obtain $U_2 = \frac{2}{7}$, $U_3 = U_4 = \frac{1}{4}$ in the same manner. Condition (C_2) holds since $\alpha_2 U^{I_4\setminus\{2\}} = U_1 + U_3 + U_4 = \frac{5}{6} < 1$. The set $[0, U^{I_4\setminus\{1\}}] \cap \gamma_1$ is a triangle with vertices $P_1 = (0, \frac{2}{7}, \frac{1}{4}, \frac{3}{14})^T$, $P_2 = (0, \frac{2}{7}, \frac{11}{56}, \frac{1}{4})^T$ and $P_3 = (0, \frac{1}{4}, \frac{1}{4}, \frac{1}{4})^T$. Since $\alpha_j P_i > 1$ for all $i \in I_3$ and $j \in I_4 \setminus \{1\}$, $[0, U^{I_4\setminus\{1\}}] \cap \gamma_1$ is above γ_j for every $j \in I_4 \setminus \{1\}$. Thus, condition (C_1) also holds. Then $\gamma_1 \cap \gamma_2 \cap \pi_3 \cap \pi_4 = \{x^*\}$ with $x^* = (\frac{1}{4}, \frac{1}{4}, 0, 0)^T$. Since $\alpha_3 x^* = 1$ and $\alpha_4 x^* = 1$, by theorem 2.5, x^* is a global attractor.

Another extreme case of theorem 2.5 when $J = \{k\}$ is simplified to the corollary below.

Corollary 2.8 *Assume that condition (C_k) and $a_{kk} = a_{jk}$ hold for some $k \in I_n$ and all $j \in I_n \setminus \{k\}$. Then $Y^{\{k\}}$ is a global attractor. The conclusion still holds if U is replaced by Y and $a_{kk} = a_{jk}$ is replaced by $a_{kk} \leq a_{jk}$.*

Remark 2.9 The conditions of corollary 2.8 with $k = 1$ and (1.4) for $i, j \in I_n$ with $i \neq j$ are mutually exclusive.

If $U_\ell = 0$ for some $\ell \in I_n$, from the definition of U we make the following observations:

(*i*) For each $j \in I_n \setminus \{\ell\}$, $Y^{\{\ell\}}$ is below γ_j and every y in γ_ℓ is on or below γ_j. Hence, $\gamma_j \cap [0, U^{I_n \setminus \{\ell\}}]$ cannot be below γ_ℓ so condition (C_ℓ) does not hold.

(*ii*) For $j \in I_n \setminus \{\ell\}$, $Y^{\{j\}} \in \gamma_j$ so, by (*i*), $Y^{\{j\}}$ cannot be below γ_ℓ. Thus, $U^{I_n \setminus \{\ell\}} = Y^{I_n \setminus \{\ell\}}$.

(*iii*) From lemma 5.4 given in §5 we shall see that every solution of (1.1) in $\mathrm{int}\mathbb{R}^n_+$ satisfies $\lim_{t\to\infty} x_\ell(t) = 0$.

Based on these observations we can redefine $U^{I_n \setminus \{\ell\}}$ by the same definition as before except the substitution of $I_n \setminus \{\ell\}$ for I_n. Repeat this process until no more new zero components occur.

Theorem 2.10 *Let $J = \{j_1, j_2, \ldots, j_m\} \subset I_n$ with $1 \le m < n$ and let V be defined by $V^J = 0$ and $V^{I_n \setminus J} \in [0, Y^{I_n \setminus J}]$ given by the definition of U with the replacement of I_n by $I_n \setminus J$. Assume that the following hold:*

(*i*) *For each $\ell \in I_m$, $Y^{\{j_\ell\}}$ is below γ_k and every y in $\gamma_{j_\ell} \cap (\pi_{j_1} \cap \cdots \cap \pi_{j_{\ell-1}})$ is on or below γ_k for all $k \in I_n \setminus \{j_1, j_2, \ldots, j_\ell\}$.*

(*ii*) *For each $k \in I_n \setminus J$, either $V^{I_n \setminus \{k\}}$ is below γ_k or $\gamma_k \cap [0, V^{I_n \setminus \{k\}}]$ is above γ_j for all $j \in I_n \setminus (J \cup \{k\})$.*

Then $(\cap_{j\in J}\pi_j) \cap (\cap_{k \in I_n \setminus J}\gamma_k) = \{x^*\}$ with $0 < x^*_k \le V_k$ for $k \in I_n \setminus J$ and x^* is a global attractor.

Remark 2.11 Suppose the conditions of theorem 2.10 are met. Then, applying theorem 2.10 to (1.1) on $\cap_{\ell \in S}\pi_\ell$ for any subset $S \subset J$, we see that every solution of (1.1) with $x_k(0) > 0$ for $k \in I_n \setminus J$ and $x_j(0) \ge 0$ for $j \in J$ satisfies $\lim_{t\to\infty} x(t) = x^*$. This remark also applies to corollary 2.13.

Example 2.12 Consider system (1.1) with $n = 5$ and

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} & a_{15} \\ a_{21} & a_{22} & a_{23} & a_{24} & a_{25} \\ a_{31} & a_{32} & a_{33} & a_{34} & a_{35} \\ a_{41} & a_{42} & a_{43} & a_{44} & a_{45} \\ a_{51} & a_{52} & a_{53} & a_{54} & a_{55} \end{pmatrix} = \begin{pmatrix} 2.1 & 1.9 & 1.9 & 0 & 1 \\ 1 & 3 & 1 & 2 & 0 \\ 1 & 1 & 3 & 1 & 1 \\ 3 & 3 & 3 & 2.1 & 0 \\ 4 & 3 & 3 & 2.5 & 2 \end{pmatrix}.$$

Since $a_{5j} \ge a_{ij}$ and $a_{55} > a_{i5}$ for all $i, j \in I_4$, $Y^{\{5\}}$ is below γ_j and each $y \in \gamma_5$ is on or below γ_j for all $j \in I_4$. Also, $a_{4j} \ge a_{ij}$ and $a_{44} > a_{i4}$ for all $i, j \in I_3$, so $Y^{\{4\}}$ is below γ_j and each $y \in \gamma_4 \cap \pi_5$ is on or below γ_j for all $j \in I_3$. Now with $J = \{j_1, j_2\}$, $j_1 = 5$ and $j_2 = 4$, condition (*i*) of theorem 2.10 is met. Since the entries a_{ij} for $i, j \in I_3$ are identical with those given in (2.4), from example 2.3 we obtain $V = (\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, 0, 0)^T$ and $\alpha_i V^{I_5 \setminus \{i\}} < 1$ for all $i \in I_5 \setminus J = I_3$. By theorem 2.10, the equilibrium $x^* = (\frac{1}{23}, \frac{11}{46}, \frac{11}{46}, 0, 0)^T$ is a global attractor.

When $n = m + 1$ with $I_n = J \cup \{k\}$, $V^{I_n \setminus \{k\}} = 0$ so assumption (*ii*) is automatically met and theorem 2.10 has the following simplified form.

Corollary 2.13 *Assume that, for each $\ell \in I_{n-1}$ and every $j \in I_n \setminus I_\ell$, $Y^{\{\ell\}}$ is below γ_j and every y in $\gamma_\ell \cap (\pi_1 \cap \cdots \cap \pi_{\ell-1})$ is on or below γ_j. Then $Y^{\{n\}}$ is a global attractor.*

Remark 2.14 The condition of corollary 2.13 is equivalent to

$$a_{j\ell} < a_{\ell\ell}, \quad a_{jj} \le a_{j-1j} \le \cdots \le a_{1j}, \quad 1 \le \ell < j \le n. \tag{2.5}$$

Note that (1.4) and (2.5) are mutually exclusive. Thus, neither of corollary 2.13 and the result given by Zeeman [14] is covered by the other.

Finally, we make a brief remark about the conditions of the main result given in [11]. It is assumed that, for some $r < n$, (1.2) with the replacement of n by r holds for all $i \in I_r$ This implies the existence of an equilibrium $x^* \in \cap_{j\in I_n\setminus I_r}\pi_j$ with $x_i^* > 0$ for $i \in I_r$. It is also assumed that x^* is above γ_j for every $j \in I_n \setminus I_r$. Then, with another condition requiring the existence of $n - r$ constants satisfying an inequality, it is shown that x^* is a global attractor. These conditions and those of theorems 2.5 and 2.10 are mutually independent though they have some overlap.

3 Analytical definition of U and algebraic expression of (C_k)

The definition of U given in §2 can be formulated as follows. For each $i \in I_n$, let

$$U_i = \begin{cases} a_{ii}^{-1} & \text{if } a_{ii} \le a_{ji} \text{ or } a_{ij} = 0 \text{ for some } j \in I_n \setminus \{i\}, \\ 0 & \text{if } a_{ji} < a_{ii}, a_{jk} \le a_{ik} \text{ for all } j,k \in I_n \setminus \{i\}, \\ \max\left\{\frac{a_{kj}-a_{ij}}{a_{ii}a_{kj}-a_{ij}a_{ki}} : j,k \in I_n \setminus \{i\}, a_{kj} > a_{ij}\right\} & \text{otherwise.} \end{cases} \tag{3.1}$$

Remark 3.1 If condition (C_k) holds for some $k \in I_n$, then every equilibrium of (1.1) on π_k is below γ_k. Indeed, $u = 0$ is obviously below γ_k. Let $u \in \pi_k$ with $u \ne 0$ be an equilibrium. Then either $u = Y^{\{i\}} \notin [0, U^{I_n\setminus\{k\}}]$ for some $i \in I_n \setminus \{k\}$ or, by the definition of U, $u \in \gamma_j \cap [0, U^{I_n\setminus\{k\}}]$ for some $j \in I_n \setminus \{k\}$. In the former case, $U_i < a_{ii}^{-1}$. By (3.1), $u = Y^{\{i\}}$ is below γ_ℓ for all $\ell \in I_n \setminus \{i\}$ so u is below γ_k. In the latter case, u is below γ_k by (C_k).

In some situation, (3.1) has a simpler form. To see this, suppose $Y^{\{i\}}$ is below γ_j for all $j \in I_n \setminus \{i\}$, i.e.

$$a_{ii} > a_{ji}, \quad j \in I_n \setminus \{i\}. \tag{3.2}$$

For each $j \in I_n \setminus \{i\}$, suppose also that either

$$a_{ik} \ge a_{jk}, \quad k \in I_n \setminus \{i\} \tag{3.3}$$

or $a_{jj} > a_{ij}$ and $\gamma_i \cap \gamma_j \cap (\cap_{\ell\in I_n\setminus\{i,j\}}\pi_\ell) = \{E^{(ij)}\}$ which is below γ_ℓ for all $\ell \in I_n \setminus \{i,j\}$. The latter can be written

$$a_{jj} > a_{ij}, \quad \frac{a_{\ell i}(a_{jj} - a_{ij}) + a_{\ell j}(a_{ii} - a_{ji})}{a_{ii}a_{jj} - a_{ij}a_{ji}} < 1, \quad \ell \in I_n \setminus \{i,j\}. \tag{3.4}$$

Then (3.1) can be simplified as

$$U_i = \max\left\{0, \frac{a_{jj} - a_{ij}}{a_{ii}a_{jj} - a_{ij}a_{ji}} : j \in I_n \setminus \{i\}, a_{jj} > a_{ij}\right\}. \tag{3.5}$$

If condition (C_k) holds for all $k \in I_n$, then, by remark 3.1, (3.2) and (3.4) hold for all $i, j \in I_n$ with $i \ne j$ so U can be defined by

$$U_i = \max\left\{\frac{a_{jj} - a_{ij}}{a_{ii}a_{jj} - a_{ij}a_{ji}} : j \in I_n \setminus \{i\}\right\}, \quad i \in I_n. \tag{3.6}$$

Lemma 3.2 *(i) For any $J \subset I_n$ with $|J| > 1$, condition (C_i) for all $i \in J$ implies*

$$\max\left\{0, \frac{a_{ij}}{a_{jj}}(1 - \alpha_j U^{I_n\setminus\{i,j\}})\right\} < 1 - \alpha_i U^{I_n\setminus\{i,j\}} \tag{3.7}$$

for all $i, j \in J$ with $i \neq j$. (ii) Conversely, for a fixed $i \in I_n$, (3.7) for all $j \in I_n \setminus \{i\}$ implies condition (C_i). (iii) Condition (C_i) holds for all $i \in I_n$ if and only if (3.7) holds for all $i, j \in I_n$ with $i \neq j$.

Proof. (i) Suppose $\alpha_i U^{I_n\setminus\{i,j\}} \geq 1$ for some $i, j \in J$ with $i \neq j$. Then $[0, U^{I_n\setminus\{i\}}] \cap \gamma_i \supset [0, U^{I_n\setminus\{i,j\}}] \cap \gamma_i \neq \emptyset$. Since $[0, U^{I_n\setminus\{i,j\}}] \cap \gamma_i$ is above γ_j by (C_i) whereas it is below γ_j by (C_j), this is impossible. Therefore, we must have

$$\alpha_i U^{I_n\setminus\{i,j\}} < 1, \quad i, j \in J, i \neq j. \tag{3.8}$$

For any $i, j \in J$ with $i \neq j$, we assume $I_n \setminus \{i, j\} \neq \emptyset$ as otherwise (3.7) holds obviously. If we can show that

$$\alpha_j U^{I_n\setminus\{i\}} \geq 1 \tag{3.9}$$

for all $j \in J \setminus \{i\}$, then (3.8) and (3.9) result in $\gamma_j \cap [U^{I_n\setminus\{i,j\}}, U^{I_n\setminus\{i\}}] = \{y\}$ with $y^{I_n\setminus\{j\}} = U^{I_n\setminus\{i,j\}}$ and $y_j = a_{jj}^{-1}(1 - \alpha_j U^{I_n\setminus\{i,j\}})$. By condition (C_i) y is below γ_i, so

$$\alpha_i y = \alpha_i U^{I_n\setminus\{i,j\}} + a_{ij}a_{jj}^{-1}(1 - \alpha_j U^{I_n\setminus\{i,j\}}) < 1$$

and (3.7) follows from this and (3.8). For any $j \in J \setminus \{i\}$, (3.9) is obvious if $U_j = a_{jj}^{-1}$. If $U_j < a_{jj}^{-1}$, then, for any $k \in I_n \setminus \{i, j\}$, $Y^{\{j\}}$ is below γ_k by the definition of U_j. On the other hand, $Y^{\{k\}}$ is below γ_j by (C_j) and remark 3.1. Consequently, $\gamma_j \cap \gamma_k \cap (\cap_{\ell \in I_n\setminus\{j,k\}} \pi_\ell) = \{E^{(jk)}\}$. Then (3.9) follows from $\alpha_j E^{(jk)} = 1$ and $E^{(jk)} \in [0, U^{I_n\setminus\{i\}}]$ by the definition of U.

(ii) If $\alpha_i U^{I_n\setminus\{i\}} < 1$ then $U^{I_n\setminus\{i\}}$ is below γ_i so condition (C_i) holds. Now suppose $\alpha_i U^{I_n\setminus\{i\}} \geq 1$. Then, for each $j \in I_n \setminus \{i\}$, since $\alpha_i U^{I_n\setminus\{i,j\}} < 1$ by (3.7), we have $U_j > 0$ and $Z^{(j)} \in [U^{I_n\setminus\{i,j\}}, U^{I_n\setminus\{i\}}]$ such that $\alpha_i Z^{(j)} = 1$. It can be shown that

$$\gamma_i \cap [0, U^{I_n\setminus\{i\}}] = \left\{ \sum_{j \in I_n\setminus\{i\}} d_j Z^{(j)} : d_j \geq 0, \sum_{j \in I_n\setminus\{i\}} d_j = 1 \right\}.$$

Hence, to show that condition (C_i) is met, we need only verify

$$\alpha_m Z^{(j)} > 1 \tag{3.10}$$

for each fixed $j \in I_n \setminus \{i\}$ and all $m \in I_n \setminus \{i\}$. Note that $Z_j^{(j)} = a_{ij}^{-1}(1 - \alpha_i U^{I_n\setminus\{i,j\}})$. It then follows from this and (3.7) that

$$\alpha_j Z^{(j)} = \alpha_j U^{I_n\setminus\{i,j\}} + a_{jj}a_{ij}^{-1}(1 - \alpha_i U^{I_n\setminus\{i,j\}}) > 1 \tag{3.11}$$

for all $j \in I_n \setminus \{i\}$. For any $m \in I_n \setminus \{i, j\}$, if $a_{mj} = 0$ then $\alpha_m Z^{(j)} = \alpha_m U^{I_n\setminus\{i\}} \geq \alpha_m Z^{(m)} > 1$ by (3.11). If $a_{mj} > 0$ and $U_m = a_{mm}^{-1}$, we also have $\alpha_m Z^{(j)} \geq a_{mm}U_m + a_{mj}Z_j^{(j)} > a_{mm}U_m = 1$. Since $a_{mj} = 0$ implies $U_m = a_{mm}^{-1}$, we have actually verified (3.10) when $U_m = a_{mm}^{-1}$. So our

remaining task is to show (3.10) under the assumption $U_m < a_{mm}^{-1}$, which implies $a_{mm} > a_{km}$ and $a_{mk} > 0$ for all $k \in I_n \setminus \{m\}$ by the definition of U_m.

(A) Suppose $I_n \neq \{i, j, m\}$. Then, for $k \in I_n \setminus \{i, j, m\}$, we have either $0 < a_{kk} \leq a_{mk}$ or $0 < a_{mk} < a_{kk}$. In the former case, $U_k = a_{kk}^{-1}$ so $\alpha_m Z^{(j)} > a_{mk} U_k \geq 1$. In the latter case, $\gamma_m \cap \gamma_k \cap (\cap_{\ell \in I_n \setminus \{m,k\}} \pi_\ell) = \{E^{(mk)}\}$. Since $E^{(mk)} \leq U^{\{m,k\}}$ and $\alpha_m E^{(mk)} = 1$, we have $\alpha_m Z^{(j)} > \alpha_m U^{\{m,k\}} \geq \alpha_m E^{(mk)} = 1$.

(B) Suppose $I_n = \{i, j, m\}$. Since $\alpha_j U^{I_n \setminus \{i\}} = a_{jj} U_j + a_{jm} U_m \geq \alpha_j Z^{(j)} > 1$ by (3.11) and $a_{jj} U_j \leq 1$, we must have $a_{jm} U_m > 0$ so $a_{mm} > a_{jm} > 0$. If $a_{mj} \geq a_{jj}$ then $\alpha_m Z^{(j)} \geq \alpha_j Z^{(j)} > 1$. Otherwise, we have $a_{jj} > a_{mj} > 0$ so $\gamma_j \cap \gamma_m \cap \pi_i = \{E^{(jm)}\}$ with

$$(E_j^{(jm)}, E_m^{(jm)}) = \left(\frac{a_{mm} - a_{jm}}{a_{jj} a_{mm} - a_{jm} a_{mj}}, \frac{a_{jj} - a_{mj}}{a_{jj} a_{mm} - a_{jm} a_{mj}} \right).$$

By the definition of U we have $E^{(jm)} \leq U^{\{j,m\}}$. Since $E_m^{(jm)} \leq U_m$ is equivalent to $a_{jj}^{-1}(1 - a_{jm} U_m) \geq a_{mj}^{-1}(1 - a_{mm} U_m)$ and (3.11) gives $a_{ij}^{-1}(1 - a_{im} U_m) > a_{jj}^{-1}(1 - a_{jm} U_m)$, from these we obtain

$$\begin{aligned} \alpha_m Z^{(j)} &= a_{mm} U_m + a_{mj} a_{ij}^{-1}(1 - a_{im} U_m) \\ &> a_{mm} U_m + a_{mj} a_{mj}^{-1}(1 - a_{mm} U_m) \\ &= 1. \end{aligned}$$

Therefore, (3.10) holds for all $m \in I_n \setminus \{i\}$.

(*iii*) This follows from (*i*) and (*ii*).

4 Restatement of the main results

With the preparation in §3 we are now in a position to restate the theorems and corollaries given in §2.

Theorem 4.1 *Assume that (1.1) satisfies (3.2), (3.4) and (3.7) for all $i, j \in I_n$ with $i \neq j$, where U is given by (3.6). Then (1.1) has an equilibrium $x^* \in \mathrm{int}\mathbb{R}^n_+$ with $x^* \leq U$ that is a global attractor.*

Corollary 4.2 *Under the conditions of theorem 4.1, for each nonempty set $J \subset I_n$, (1.1) has an equilibrium $x_J \in [0, U^J]$ with $(x_J)_i > 0$ for $i \in J$ that is a global attractor within the interior of $\cap_{\ell \in I_n \setminus J} \pi_\ell$ viewed as $\mathbb{R}_+^{|J|}$. Moreover, (1.1) has 2^n equilibria in $\mathbb{R}^n_+$.*

From lemma 3.2 we see that the conditions of theorem 4.3 and corollary 4.4 given below are slightly stronger than those of theorem 2.5 and corollary 2.8 respectively.

Theorem 4.3 *Let $J \subset I_n$ with $J \neq \emptyset$. Assume that (3.7) holds for each $i \in J$ and every $j \in I_n \setminus \{i\}$, where each component U_i of U is defined either by (3.1) or, if (3.2) and (3.4) holds for all $j \in I_n \setminus \{i\}$, by (3.6). Then (1.1) has an equilibrium $x^* \in [0, U^J]$ with $x_j^* > 0$ for $j \in J$. In addition, if $\alpha_j x^* = 1$ for all $j \in I_n \setminus J$, then x^* is a global attractor. Moreover, "$\alpha_j x^* = 1$" can be relaxed to "$\alpha_j x^* \geq 1$" when U is replaced by Y given by (2.1).*

Corollary 4.4 *Assume that (3.7) and $a_{ii} = a_{ji}$ hold for some $i \in I_n$ and all $j \in I_n \setminus \{i\}$, where U is defined by (3.1). Then $Y^{\{i\}}$ is a global attractor of (1.1). Moreover, "$a_{ii} = a_{ji}$" can be replaced by "$a_{ii} \le a_{ji}$" if U is replaced by Y.*

Theorem 4.5 *Let $J = \{j_1, j_2, \dots, j_m\} \subset I_n$ with $1 \le m < n$ and let $U \in \mathbb{R}^n_+$ be defined by $U_i = 0$ for $i \in J$ and by (3.1), with I_n replaced by $I_n \setminus J$, for $i \in I_n \setminus J$. Assume that*

(i) for each $\ell \in I_m$ and every $j, k \in I_n \setminus \{j_1, j_2, \dots, j_\ell\}$,

$$a_{kj_\ell} < a_{j_\ell j_\ell}, \quad a_{kj} \le a_{j_\ell j}; \tag{4.1}$$

(ii) for all $i, j \in I_n \setminus J$ with $i \ne j$ (if $n > m+1$),

$$\max\left\{0, \frac{a_{ij}}{a_{jj}}(1 - \alpha_j U^{(I_n \setminus J)\setminus\{i,j\}})\right\} < 1 - \alpha_i U^{(I_n \setminus J)\setminus\{i,j\}}. \tag{4.2}$$

Then (1.1) has an equilibrium $x^* \in [0, U^{I_n \setminus J}]$ with $x^*_j > 0$ for $j \in I_n \setminus J$ that is a global attractor.

Corollary 4.6 *Assume that (1.1) satisfies (2.5). Then $Y^{\{n\}}$ is a global attractor.*

Remark 4.7 By remark 3.1 and theorem 2.1, we are tempted to make the following conjecture as an improvement of theorem 2.1:

> System (1.1) has an equilibrium $x^* \in \mathrm{int}\mathbb{R}^n_+$ that is a global attractor if, for each $i \in I_n$, every equilibrium of (1.1) on π_i is below γ_i.

This is obviously true when $n = 2$. When $n = 3$, the condition ensures that each $Y^{\{i\}}$ is a local repellor and, by lemma 5.1 given in the next section, $\gamma_1 \cap \gamma_2 \cap \gamma_3 = \{x^*\} \subset \mathrm{int}\mathbb{R}^3_+$. Then, by a result given in [4], x^* is a global attractor. Hence, it is fairly reasonable to make the above conjecture for general n-dimensional system (1.1). However, further investigation is needed to clarify it.

5 The proofs of theorems 2.5 and 2.10

The method of ultimate contracting cells (see [8, Theorem 2] or [9, Lemma 1]) will be used in the proofs of the theorems though it is not explicitly stated here. Our first lemma establishes the existence of a unique equilibrium $x^* \in \mathrm{int}\mathbb{R}^n_+$ under conditions weaker than (C_k) for all $k \in I_n$.

Lemma 5.1 *Assume that, for each $i \in I_n$, every equilibrium of (1.1) on π_i is below γ_i. Then there is a unique equilibrium $x^* \in \mathrm{int}\mathbb{R}^n_+$ with $x^* \le Y$.*

Proof. Denoting the statement of lemma 5.1 by $P(n)$, we show the truth of $P(n)$ by induction. The truth of $P(n)$ is trivial for $n = 1$ and $n = 2$. Supposing that $P(n)$ is true for some $k \ge 2$ and all $n \in I_k$, we now show the truth of $P(k+1)$.

By the assumption, for each $i \in I_{k+1}$, every equilibrium of (1.1) on π_i is below γ_i. In particular, for each $i \in I_k$, every equilibrium of (1.1) on $\pi_i \cap \pi_{k+1}$ is below γ_i and γ_{k+1}. Viewing (1.1) on π_{k+1} as a k-dimensional system on $\mathbb{R}^k_+$ and using the truth of $P(k)$, we have a unique




Figure 5.1

equilibrium u^* of (1.1) on π_{k+1} satisfying $0 < u_i^* \le a_{ii}^{-1}$ for $i \in I_k$. This implies that the solution of the system

$$\alpha_i u = 1, \quad i \in I_k$$

can be written $u(\delta)$ with $u(0) = u^*$, $u_{k+1}(\delta) = \delta$ and $u(\delta) - u^*$ is linear in δ. Thus, $u(\delta) \in \text{int}\mathbb{R}_+^{k+1}$ holds for sufficiently small $\delta > 0$.

(i) If $u(a_{k+1k+1}^{-1}) \in \text{int}\mathbb{R}_+^{k+1}$ then $u(\delta) \in \text{int}\mathbb{R}_+^{k+1}$ for all $\delta \in (0, a_{k+1k+1}^{-1}]$ and

$$\alpha_{k+1} u(a_{k+1k+1}^{-1}) \ge a_{k+1k+1} a_{k+1k+1}^{-1} = 1.$$

As $\alpha_{k+1} u(0) = \alpha_{k+1} u^* < 1$ by the assumption, there is a $\delta_0 \in (0, a_{k+1k+1}^{-1}]$ such that $\alpha_{k+1} u(\delta_0) = 1$. Then $u(\delta_0) \in \text{int}\mathbb{R}_+^{k+1}$ is the required equilibrium.

(ii) If $u(a_{k+1k+1}^{-1}) \notin \text{int}\mathbb{R}_+^{k+1}$, then there is a $\delta_1 \in (0, a_{k+1k+1}^{-1}]$ such that $u(\delta) \in \text{int}\mathbb{R}_+^{k+1}$ for all $\delta \in (0, \delta_1)$ but $u_\ell(\delta_1) = 0$ for some $\ell \in I_k$. Thus $\cap_{i \in I_k} \gamma_i$ is a line segment joining u^* with $u(\delta_1)$. Again, by the assumption and the truth of $P(k-1)$ and $P(k)$, (1.1) has a unique equilibrium v^* on $\pi_\ell \cap \pi_{k+1}$ satisfying $0 < v_i^* \le a_{ii}^{-1}$ for $i \in I_k \setminus \{\ell\}$ and a unique equilibrium w^* on π_ℓ with $0 < w_j^* \le a_{jj}^{-1}$ for $j \in I_{k+1} \setminus \{\ell\}$. Consequently, the set $S = \cap_{i \in I_k \setminus \{\ell\}} \gamma_i$ is a two-dimensional plane and $\cap_{i \in I_{k+1} \setminus \{\ell\}} \gamma_i$ is a line segment starting from w^* on S (see Figure 5.1). Note that $S \cap \pi_\ell$ is a line segment containing v^*, w^* and $u(\delta_1)$, and $S \cap \pi_{k+1}$ is a line segment containing v^* and u^*. Since v^* is below both γ_ℓ and γ_{k+1}, u^* is below γ_{k+1} and w^* is below γ_ℓ, we must have $\gamma_\ell \cap \gamma_{k+1} \cap S = \{x^*\} \subset \text{int}\mathbb{R}_+^n$ with x^* the required equilibrium.

Therefore, we have shown the truth of $P(k+1)$ when $P(n)$ is true for all $n \in I_k$. By induction, $P(n)$ is true for all $n \ge 1$.

The next lemma gives an ultimate upper bound of any solution of (1.1) in terms of a given ultimate lower bound (see [7, Lemma 3.2]).

Lemma 5.2 *Let $\hat{x} \in \mathbb{R}_+^n$ satisfy $\hat{x}_i = 0$ if $\alpha_i \hat{x} > 1$ for any $i \in I_n$ and let $\hat{y} \in \mathbb{R}_+^n$ be given by*

$$\hat{y}_j = \max\{0, \hat{x}_j + a_{jj}^{-1}(1 - \alpha_j \hat{x})\}, \quad j \in I_n. \tag{5.1}$$

If a solution of (1.1) in $\text{int}\mathbb{R}_+^n$ satisfies $\underline{\lim}_{t\to\infty} x(t) \ge \hat{x}$, then it also satisfies $\overline{\lim}_{t\to\infty} x(t) \le \hat{y}$.

However, we shall need a refined ultimate upper bound based on (5.1). For the above $\hat{x}$ and some $j \in I_n$, if we know that $x_j(t) \to \hat{x}_j$ as $t \to \infty$, then we take $\hat{y}_j = \hat{x}_j$ rather than (5.1). Now for $\hat{y} \ge \hat{x}$ with $\hat{y}_j = \hat{x}_j$ or $\hat{y}_j$ given by (5.1) for each $j \in I_n$, let $J_0 = \{i \in I_n : \hat{x}_i < \hat{y}_i\}$ and

define $\hat{z} \in [\hat{x}, \hat{y}]$ by (5.2) for each $i \in I_n$,

$$\hat{z}_i = \begin{cases} \hat{y}_i & \text{if } J_0 = \{i\} \text{ or } i \notin J_0 \text{ or, for some } j \in J_0 \setminus \{i\}, \\ & a_{ij} = 0 \text{ or } \alpha_j \hat{x}^{I_n \setminus \{i\}} + a_{ji}\hat{y}_i \geq 1; \\ \hat{x}_i & \text{if } i \in J_0 \neq \{i\}, \alpha_j \hat{x}^{I_n \setminus \{i\}} + a_{ji}\hat{y}_i < 1 \text{ and } \alpha_j y \leq 1 \\ & \text{for all } y \in \gamma_i \cap [\hat{x}, \hat{y}] \text{ and } j \in J_0 \setminus \{i\}; \\ \max\{u_i^{(ijk)} : \gamma_i \cap \gamma_k \cap [\hat{x}, H^{(ij)}] = \{u^{(ijk)}\}, j, k \in J_0 \setminus \{i\}\} & \\ & \text{otherwise,} \end{cases} \tag{5.2}$$

where $H^{(ij)} = \hat{x}^{I_n \setminus \{i,j\}} + \hat{y}^{\{i,j\}}$.

Lemma 5.3 *Assume that a solution $x(t)$ of (1.1) in $\mathrm{int}\mathbb{R}^n_+$ satisfies*

$$\hat{x} \leq \underline{\lim}_{t\to\infty} x(t) \leq \overline{\lim}_{t\to\infty} x(t) \leq \hat{y}$$

for some $\hat{x}, \hat{y} \in \mathrm{int}\mathbb{R}^n_+$, where, for any $i, j \in I_n$, $\hat{x}_i = 0$ if $\alpha_i \hat{x} > 1$ and $\hat{y}_j$ is given by (5.1) if $\hat{y}_j > \hat{x}_j$. Then, for $\hat{z} \geq \hat{y}$ defined by (5.2), $x(t)$ also satisfies $\overline{\lim}_{t\to\infty} x(t) \leq \hat{z}$.

Proof. We only need to show $\overline{\lim}_{t\to\infty} x_i(t) \leq \hat{z}_i$ if $\hat{z}_i < \hat{y}_i$ for a fixed $i \in I_n$. This can be achieved by showing that, for every $r_0 \in (\hat{z}_i, \hat{y}_i]$, there is an $r_1 \in [\hat{z}_i, r_0)$ such that

$$\overline{\lim}_{t\to\infty} x_i(t) \leq r_0 \tag{5.3}$$

implies

$$\overline{\lim}_{t\to\infty} x_i(t) \leq r_1. \tag{5.4}$$

From $\hat{z}_i < \hat{y}_i$ and (5.2) we know that $i \in J_0 \neq \{i\}$, $a_{ij} > 0$ and $\alpha_j \hat{x}^{I_n \setminus \{i\}} + a_{ji}\hat{y}_i < 1$, i.e. $\hat{x}^{I_n \setminus \{i\}} + \hat{y}^{\{i\}}$ is below γ_j, for all $j \in J_0 \setminus \{i\}$. By (5.1), $\hat{x}^{I_n \setminus \{k\}} + \hat{y}^{\{k\}}$ is on γ_k for all $k \in J_0$. Thus, for each $j \in J_0 \setminus \{i\}$, $H^{(ij)} = \hat{x}^{I_n \setminus \{i,j\}} + \hat{y}^{\{i,j\}}$ is on or above γ_j and $\hat{x}^{I_n \setminus \{i\}} + \hat{y}^{\{i\}}$ is below γ_k for every $k \in J_0 \setminus \{i\}$. Now suppose (5.3) holds. The line segment $L_{r_0} = \{x \in [\hat{x}, H^{(ij)}] : x_i = r_0\}$ must intersect a plane γ_k for some $k \in J_0 \setminus \{i\}$ at $Q^{(j)}$ (see Figure 5.2) such that $Q^{(j)}_i = r_0$, $Q^{(j)}_\ell = \hat{x}_\ell$ for $\ell \in I_n \setminus \{i, j\}$ and

$$Q^{(j)}_j = \min\{a^{-1}_{\ell j}(1 - \alpha_\ell \hat{x}^{I_n \setminus \{i,j\}} - a_{\ell i} r_0) : \ell \in J_0 \setminus \{i\}, a_{\ell j} \neq 0\}.$$

By the definition of $\hat{z}_i$ and $r_0 > \hat{z}_i$, we see that $Q^{(j)}_j > \hat{x}_j$ and $Q^{(j)}$ is above γ_i but on or below γ_ℓ for every $\ell \in J_0 \setminus \{i\}$. Put $\gamma(\varepsilon) = \{x \in \mathbb{R}^n_+ : \beta x = \varepsilon\}$, where $\beta = (\beta_1, \beta_2, \ldots, \beta_n)$ with $\beta_j = 0$ for $j \in \{i\} \cup (I_n \setminus J_0)$ and

$$\beta_j = \frac{(Q^{(j)}_j - \hat{x}_j)^{-1}}{1 + \sum_{\ell \in J_0 \setminus \{i\}} (Q^{(\ell)}_\ell - \hat{x}_\ell)^{-1} \hat{x}_\ell}$$

for $j \in J_0 \setminus \{i\}$. Then the x_i-axis is below and parallel to the plane $\gamma(\varepsilon)$ for each $\varepsilon > 0$, $[\hat{x}, \hat{x}^{I_n \setminus \{i\}} + \hat{y}^{\{i\}}]$ is below and parallel to $\gamma(1)$, and $Q^{(j)}$ is on $\gamma(1)$ for $j \in J_0 \setminus \{i\}$. Moreover, for any $\varepsilon \in (\beta\hat{x}, 1)$, the set

$$A(\varepsilon) = \{x \in [\hat{x}, \hat{y}] : x_i \leq r_0, \beta x \leq \varepsilon\} \tag{5.5}$$




Figure 5.2

is below γ_j for every $j\in J_0\setminus\{i\}$. We now show the inequality

$$\underline{\lim}_{t\to\infty}\beta x(t)\geq 1 \tag{5.6}$$

by assuming its falsity, i.e. $\underline{\lim}_{t\to\infty}\beta x(t)=\delta$ for some $\delta<1$. Then there is a sequence $\{t_m\}$ satisfying $t_m\to\infty$ as $m\to\infty$ and $\beta x(t_m)<(1+\delta)/2$ for all $m\geq 1$. Since $\lim_{t\to\infty}x_\ell(t)=\hat{x}_\ell$ for $\ell\in I_n\setminus J_0$, (5.3) holds and $A(\frac{1+\delta}{2})$ given by (5.5) is a closed set, we can assume the existence of $T>0$, $[\tilde{x},\tilde{y}]\supseteq[\hat{x},\hat{y}]$ and $\tilde{r}_0>r_0$ such that $x(t)\in[\tilde{x},\tilde{y}]$ and $x_i(t)\leq\tilde{r}_0$ for all $t\geq T$ and the set

$$\tilde{A}((1+\delta)/2)=\{x\in[\tilde{x},\tilde{y}]:x_i\leq\tilde{r}_0,\beta x\leq(1+\delta)/2\}$$

is below γ_j for all $j\in J_0\setminus\{i\}$. Thus, for any $t\geq T$, $x(t)\in\tilde{A}(\frac{1+\delta}{2})$ implies $\beta x'(t)>0$. Without loss of generality, we may also assume that $x(t_m)\in\tilde{A}(\frac{1+\delta}{2})$, so that $\beta x'(t_m)>0$, for all $m\geq 1$. Then, for each $m\geq 1$, $\beta x(t)$ is increasing for $t\geq t_m$ as long as $x(t)\in\tilde{A}(\frac{1+\delta}{2})$. This, together with $\beta x(t_m)<(1+\delta)/2$ for all $m\geq 1$, results in $x(t)\in\tilde{A}(\frac{1+\delta}{2})$ for all $t\geq t_1$. By the closeness of $\tilde{A}(\frac{1+\delta}{2})$ there is an $\varepsilon\in(0,1)$ such that $\alpha_j x\leq\varepsilon$ for all $x\in\tilde{A}(\frac{1+\delta}{2})$ and $j\in J_0\setminus\{i\}$. Hence, from (1.1),

$$x_j(t)\geq x_j(t_1)e^{b_j(1-\varepsilon)(t-t_1)}$$

for $t\geq t_1$ and $j\in J_0\setminus\{i\}$. This leads to $\beta x(t)\to\infty$ as $t\to\infty$, a contradiction to $\underline{\lim}_{t\to\infty}\beta x(t)=\delta$. Therefore, we must have (5.6).

Note that, for each $j\in J_0\setminus\{i\}$, $\gamma(1)\cap[\hat{x},H^{(ij)}]$ is a line segment parallel to $[\hat{x},\hat{x}^{I_n\setminus\{i\}}+\hat{y}^{\{i\}}]$ and $Q^{(j)}$ is on it. So we let $\gamma_i\cap\gamma(1)\cap[\hat{x},H^{(ij)}]=\{P^{(j)}\}$ or, if it is empty, let $\gamma(1)\cap[\hat{x},\hat{x}^{I_n\setminus\{j\}}+\hat{y}^{\{j\}}]=\{P^{(j)}\}$, and let

$$r_1=\max\{P_i^{(j)}:j\in J_0\setminus\{i\}\}.$$

Then $r_1\in[\hat{z}_i,r_0)$ as each $Q^{(j)}$ is above γ_i. We observe that the closed set $S(\eta)=\{x\in[\hat{x},\hat{y}]:x_i\geq\eta,\beta x\geq 1\}$ is above γ_i for every $\eta>r_1$. Then, from this observation and (5.6) we see that $x(t)$ satisfies $x_i(t)<\eta$ for each $\eta>r_1$ and all sufficiently large t. Thus (5.4) holds.

From lemmas 5.2 and 5.3 we obtain an ultimate upper bound for all solutions of (1.1) in $\mathrm{int}\mathbb{R}^n_+$.

Lemma 5.4 *Every solution of (1.1) in* $\mathrm{int}\mathbb{R}^n_+$ *satisfies*

$$\overline{\lim}_{t\to\infty} x(t) \le U$$

for U *given by (3.1). If* $U_{j_1} = 0$ *for some* $j_1 \in I_n$*, then there exists* $J = \{j_1, j_2, \dots, j_m\} \subset I_n$ *with* $1 \le m < n$ *such that (4.1) holds for each* $\ell \in I_m$ *and every* $j, k \in I_n \setminus \{j_1, j_2, \dots, j_\ell\}$*. Moreover, every solution of (1.1) in* $\mathrm{int}\mathbb{R}^n_+$ *satisfies*

$$\overline{\lim}_{t\to\infty} x(t) \le V,$$

where $V_k = 0$ *for* $k \in J$ *and* $V_i \in (0, a_{ii}^{-1}]$ *is given by (3.1) with the replacement of* I_n *by* $I_n \setminus J$ *for* $i \in I_n \setminus J$*.*

Proof. By lemma 5.2, every solution of (1.1) in $\mathrm{int}\mathbb{R}^n_+$ satisfies

$$0 = \hat{x} \le \underline{\lim}_{t\to\infty} x(t) \le \overline{\lim}_{t\to\infty} x(t) \le \hat{y} = Y.$$

Note that $\hat{z}$ given by (5.2) with $[\hat{x}, \hat{y}] = [0, Y]$ coincides with U defined by (3.1). Then, by lemma 5.3, every solution of (1.1) in $\mathrm{int}\mathbb{R}^n_+$ satisfies

$$0 = \hat{x} \le \underline{\lim}_{t\to\infty} x(t) \le \overline{\lim}_{t\to\infty} x(t) \le \hat{z} \tag{5.7}$$

with $\hat{z} = U$. If $U_{j_1} = 0$ for some $j_1 \in I_n$, then, with $J = \{j_1\}$, (4.1) follows from (3.1) for $\ell = 1$ and all $j, k \in I_n \setminus \{j_1\}$. From the observation (ii) made before theorem 2.10 we know that $U = Y^{I_n \setminus \{J_1\}}$. Applying lemma 5.3 again with $[\hat{x}, \hat{y}] = [0, Y^{I_n \setminus \{j_1\}}]$, we see that every solution of (1.1) in $\mathrm{int}\mathbb{R}^n_+$ satisfies (5.7) with $\hat{z}$ given by (5.2). Note that, for each $j \in I_n \setminus \{j_1\}$, $\hat{z}_j$ coincides with U_j given by (3.1) after the replacement of I_n by $I_n \setminus \{j_1\}$. If $\hat{z}_j > 0$ for all $j \ne j_1$, then the conclusion follows with $J = \{j_1\}$ and $V = \hat{z}$. Otherwise, if $\hat{z}_{j_2} = U_{j_2} = 0$ for some $j_2 \in I_n \setminus \{j_1\}$, the conclusion follows from repeating the above process a number of times.

The next lemma gives a refined ultimate lower bound based on the given ultimate lower and upper bounds. In particular, it confirms that condition (C_k) guarantees the survival of the kth species.

Lemma 5.5 *Assume that every solution of (1.1) in* $\mathrm{int}\mathbb{R}^n_+$ *satisfies*

$$\hat{x} \le \underline{\lim}_{t\to\infty} x(t) \le \overline{\lim}_{t\to\infty} x(t) \le \hat{u} \tag{5.8}$$

for some $\hat{x}, \hat{u} \in \mathbb{R}^n_+$*. Assume also that, for some* $i \in J_1 = \{j \in I_n : \hat{u}_j > \hat{x}_j\} \ne \emptyset$*, either* (A) $\hat{x}^{\{i\}} + \hat{u}^{I_n \setminus \{i\}}$ *is below* γ_i *or* (B) $\hat{x}$ *is below* γ_i *but* $\gamma_i \cap [\hat{x}, \hat{x}^{\{i\}} + \hat{u}^{I_n \setminus \{i\}}]$ *is above* γ_j *for every* $j \in J_1 \setminus \{i\}$*. Then there is a* $\delta_0 > 0$ *such that*

$$\underline{\lim}_{t\to\infty} x_i(t) \ge \hat{x}_i + \delta_0. \tag{5.9}$$

Proof. Under (A) and (5.8) we show that

$$\delta_0 = a_{ii}^{-1}(1 - a_{ii}\hat{x}_i - \alpha_i \hat{u}^{I_n \setminus \{i\}})$$

meets the requirement of (5.9). In fact, $\delta_0 > 0$ since $\hat{x}^{\{i\}} + \hat{u}^{I_n \setminus \{i\}}$ is below γ_i. Moreover, for arbitrary $\delta \in (0, \delta_0)$, we have $\alpha_i \hat{u}^{I_n \setminus \{i\}} + a_{ii}(\delta_0 + \hat{x}_i) = 1$ and $\alpha_i \hat{u}^{I_n \setminus \{i\}} + a_{ii}(\delta + \hat{x}_i) < 1$. For each solution of (1.1) in $\mathrm{int}\mathbb{R}^n_+$, since $\overline{\lim}_{t\to\infty} x(t)^{I_n \setminus \{i\}} \le \hat{u}^{I_n \setminus \{i\}}$ by (5.8), there is a $t_0 > 0$ and

$\varepsilon \in (0,1)$ such that $\alpha_i x(t)^{I_n\setminus\{i\}} + a_{ii}(\delta + \hat{x}_i) \le \varepsilon < 1$ for all $t \ge t_0$. If $x_i(t) \le \hat{x}_i + \delta$ holds for all $t \ge t_0$ then

$$\alpha_i x(t) \le \alpha_i x(t)^{I_n\setminus\{i\}} + a_{ii}(\delta + \hat{x}_i) \le \varepsilon < 1$$

for $t \ge t_0$, so $x_i(t) \ge x_i(t_0)e^{b_i(1-\varepsilon)(t-t_0)} \to \infty$ as $t \to \infty$. This is impossible. Since $x_i(\bar{t}) \le \hat{x}_i + \delta$ for any $\bar{t} \ge t_0$ implies the increase of x_i for t at the vicinity of $\bar{t}$, there must be a $t_1 > t_0$ such that $x_i(t) > \hat{x}_i + \delta$ for all $t \ge t_1$. Thus, $\underline{\lim}_{t\to\infty} x_i(t) \ge \hat{x}_i + \delta$. Then (5.9) follows from this and the arbitrariness of $\delta \in (0, \delta_0)$.

Under (5.8) and (B), the proof of [7, Lemma 3.3] can be copied to here almost verbatim after the replacement of $[\hat{x}, \hat{y}]$ by $[\hat{x}, \hat{u}]$.

With the above preparation we are now able to embark on the proofs of the theorems.

Proof of theorem 2.5. For $k \in J$, since condition (C_k) holds, by remark 3.1 every equilibrium of (1.1) on $\pi_k \cap (\cap_{j\in I_n\setminus J}\pi_j)$ is below γ_k. Restricting (1.1) to $\cap_{j\in I_n\setminus J}\pi_j$ and applying lemma 5.1 to it, we obtain a unique equilibrium x^* on $\cap_{j\in I_n\setminus J}\pi_j$ satisfying $0 < x_k^* \le a_{kk}^{-1}$ for $k \in J$. By lemma 5.4, every solution of (1.1) in $\mathrm{int}\mathbb{R}^n_+$ satisfies

$$0 \le \underline{\lim}_{t\to\infty} x(t) \le \overline{\lim}_{t\to\infty} x(t) \le U \le Y. \tag{5.10}$$

Since condition (C_k) holds for all $k \in J$, by lemma 5.5 there is a $\delta \in (0,1]$ such that every solution of (1.1) in $\mathrm{int}\mathbb{R}^n_+$ satisfies $\underline{\lim}_{t\to\infty} x(t) \ge \delta x^*$. Let δ_0 be the supremum of such δ. Then, by lemma 5.2, every solution of (1.1) in $\mathrm{int}\mathbb{R}^n_+$ satisfies

$$\delta_0 x^* = \hat{x} \le \underline{\lim}_{t\to\infty} x(t) \le \overline{\lim}_{t\to\infty} x(t) \le \hat{y},$$

where $\hat{y}$ is given by (5.1). By the assumption, x^* is on γ_j for every $j \in I_n$. Hence, $\hat{x} = x^* = \hat{y}$, so x^* is a global attractor, if $\delta_0 = 1$.

Now suppose $\delta_0 < 1$. By lemma 5.3, every solution of (1.1) in $\mathrm{int}\mathbb{R}^n_+$ satisfies

$$\hat{x} \le \underline{\lim}_{t\to\infty} x(t) \le \overline{\lim}_{t\to\infty} x(t) \le \hat{z},$$

where $\hat{z}$ is given by (5.2). For each $k \in J$, since $\alpha_k \hat{x} = \delta_0 \alpha_k x^* = \delta_0 < 1$, if we can show that

$$\alpha_j y > 1, \quad j \in I_n \setminus \{k\} \tag{5.11}$$

whenever $y \in [\hat{x}, \hat{x}^{\{k\}} + \hat{z}^{I_n\setminus\{k\}}] \cap \gamma_k \ne \emptyset$, by lemma 5.5 again we have $\underline{\lim}_{t\to\infty} x(t) \ge \delta x^*$ for some $\delta \in (\delta_0, 1]$, which contradicts the definition of δ_0. This contradiction shows that $\delta_0 = 1$.

To prove (5.11) we introduce an affine mapping $f : [0, Y] \to [0, Y]$ defined by $f(x) = \hat{x} + (1 - \delta_0)x$. Then $f(0) = \hat{x}$ and, for each $i \in I_n$, $f_i(Y) = \hat{x}_i + (1-\delta_0)Y_i = \hat{x}_i + (1 - \alpha_i\hat{x})a_{ii}^{-1} = \hat{y}_i > \hat{x}_i$. Thus, $f([0, Y]) = [\hat{x}, \hat{y}]$ and, for any $S \subset I_n$,

$$f([0, Y^S]) = [\hat{x}, \hat{x}^{I_n\setminus S} + \hat{y}^S].$$

Next we check that $f(U) = \hat{z}$ so that $F([0, U]) = [\hat{x}, \hat{z}]$ and, for any $S \subset I_n$,

$$f([0, U^S]) = [\hat{x}, \hat{x}^{I_n\setminus S} + \hat{z}^S]. \tag{5.12}$$

For this purpose, we note that $x^* \in \cap_{i\in I_n}\gamma_i$ by the assumption and $f(x^*) = \hat{x} + (1-\delta_0)x^* = x^*$. For each $y \in [0, Y] \setminus \{x^*\}$, $f(y) = \delta_0 x^* + (1 - \delta_0)y$ is on the line segment $\overline{yx^*}$. Thus, $f(\gamma_i \cap [0, Y]) = \gamma_i \cap [\hat{x}, \hat{y}]$ for all $i \in I_n$. For any fixed $i \in I_n$, if $U_i = a_{ii}^{-1}$ then, by (3.1), $a_{ij} = 0$

or $Y^{\{i\}}$ is not below γ_j, so $f(Y^{\{i\}}) = \hat{x} + (1-\delta_0)Y^{\{i\}} = \hat{x}^{I_n\setminus\{i\}} + \hat{y}^{\{i\}}$ is not below γ_j, for some $j \in I_n \setminus \{i\}$. By (5.2) we have $\hat{z}_i = \hat{y}_i$. Therefore, $f_i(U) = f_i(Y^{\{i\}}) = \hat{y}_i = \hat{z}_i$. If $U_i = 0$, then $Y^{\{i\}}$ is below γ_j and every $y \in \gamma_i$ is on or below γ_j for all $j \in I_n \setminus \{i\}$. So $\hat{x}^{I_n\setminus\{i\}} + \hat{y}^{\{i\}}$, which is on γ_i, is below γ_j and every $y \in \gamma_i \cap [\hat{x}, \hat{y}]$ is on or below γ_j for all $j \in I_n \setminus \{i\}$. By (5.2) again, $\hat{z}_i = \hat{x}_i$. Hence, $f_i(U) = f_i(0) = \hat{x}_i = \hat{z}_i$. Now suppose $0 < U_i < a_{ii}^{-1}$. From (3.1) we have $a_{ii} > a_{ki}$ and $a_{ij} < a_{kj}$ for some $k, j \in I_n \setminus \{i\}$. Then $\gamma_i \cap \gamma_k \cap [0, Y^{\{i,j\}}]$ consists of a single point having

$$\frac{a_{kj} - a_{ij}}{a_{ii}a_{kj} - a_{ij}a_{ki}}$$

as its ith coordinate. Since $x \in \gamma_i \cap \gamma_k \cap [0, Y^{\{i,j\}}]$ if and only if $f(x) = y \in \gamma_i \cap \gamma_k \cap [\hat{x}, \hat{x}^{I_n\setminus\{i,j\}} + \hat{y}^{\{i,j\}}]$ and $\alpha_i\hat{x} = \delta_0 = \alpha_k\hat{x}$, by a routine check we see that $\gamma_i \cap \gamma_k \cap [\hat{x}, \hat{x}^{I_n\setminus\{i,j\}} + \hat{y}^{\{i,j\}}]$ also consists of a single point $u^{(ijk)}$ with

$$u_i^{(ijk)} = \hat{x}_i + (1-\delta_0)\frac{a_{kj} - a_{ij}}{a_{ii}a_{kj} - a_{ij}a_{ki}}.$$

From (3.1) and (5.2) we obtain

$$\begin{aligned} f_i(U) &= \hat{x}_i + (1-\delta_0)\max\left\{\frac{a_{kj} - a_{ij}}{a_{ii}a_{kj} - a_{ij}a_{ki}} : k, j \in I_n \setminus \{i\}, a_{kj} > a_{ij}\right\} \\ &= \max\{u_i^{(ijk)} : k, j \in I_n \setminus \{i\}, a_{kj} > a_{ij}\} \\ &= \hat{z}_i. \end{aligned}$$

Therefore, $f(U) = \hat{z}$. Now that (5.12) is derived from $f(U) = \hat{z}$, we must have

$$f([0, U^{I_n\setminus\{k\}}] \cap \gamma_k) = [\hat{x}, \hat{x}^{\{k\}} + \hat{z}^{I_n\setminus\{k\}}] \cap \gamma_k.$$

Then, for $y \in [\hat{x}, \hat{x}^{\{k\}} + \hat{z}^{I_n\setminus\{k\}}] \cap \gamma_k \neq \emptyset$, there is a unique $x \in [0, U^{I_n\setminus\{k\}}] \cap \gamma_k$ such that $f(x) = y$. By condition (C_k) for $k \in J$, x is above γ_j for all $j \in I_n \setminus \{k\}$. Thus y is above γ_j for all $j \in I_n \setminus \{k\}$, i.e. y satisfies (5.11).

If Y replaces U in the conditions and $\alpha_j x^* \geq 1$ is assumed for all $j \in I_n \setminus J$, then $f(Y) \geq \hat{y}$ so $f([0, Y]) \supset [\hat{x}, \hat{y}]$. The above argument is still valid after minor modifications.

Finally, that $x^* \leq U$ follows from (5.10) and $\lim_{t\to\infty} x(t) = x^*$ for every solution of (1.1) in $\mathrm{int}\mathbb{R}^n_+$.

Proof of theorem 2.10. When $n \geq m+2$, by assumption (ii) and theorem 2.1 with the replacement of I_n by $I_n \setminus J$ we know that (1.1) has a unique equilibrium $x^* \in \cap_{j\in J}\pi_j$ with $0 < x_i^* \leq V_i$ for $i \in I_n \setminus J$. This is also true with $x^* = Y^{\{k\}}$ when $I_n = J \cup \{k\}$ (i.e. $n = m+1$). We next show that x^* is a global attractor. From lemma 5.4 we know that every solution of (1.1) in $\mathrm{int}\mathbb{R}^n_+$ satisfies

$$0 = \hat{x} \leq \underline{\lim}_{t\to\infty} x(t) \leq \overline{\lim}_{t\to\infty} x(t) \leq \hat{z}$$

with $\hat{z} = V$. Then, by (i) and (ii), for each $k \in I_n \setminus J$, either $V^{I_n\setminus\{k\}}$ is below γ_k or $\gamma_k \cap [0, V^{I_n\setminus\{k\}}]$ is above γ_j for every $j \in I_n \setminus \{k\}$. Applying lemma 5.5 to (1.1) we obtain

$$\lim_{t\to\infty} x(t)^J = 0, \quad \underline{\lim}_{t\to\infty} x_j(t) \geq \delta, \quad j \in I_n \setminus J$$

for some $\delta > 0$ and all solutions of (1.1) in $\mathrm{int}\mathbb{R}^n_+$. Since $\{x^*\}$ is the largest invariant set of (1.1) in $\{y \in \mathbb{R}^n_+ : y^J = 0, y_j \geq \delta, j \in I_n \setminus J\}$, x^* must be a global attractor.

References

[1] S. Ahmad and A. C. Lazer, One species extinction in an autonomous competitive model, Proc. First World Congress on Nonlinear Analysis, Walter DeGruyter, Berlin, 359–368 (1995).

[2] S. Ahmad and A. C. Lazer, Necessary and sufficient average growth in a Lotka-Volterra system, Nonlinear Analysis **34**, 191–228 (1998).

[3] S. Ahmad and A. C. Lazer, Average growth and extinction in a competitive Lotka-Volterra system, Nonlinear Analysis **62**, 545–557 (2005).

[4] P. van den Driessche abd M. L. Zeeman, Three-dimensional competitive Lotka-Volterra systems with no periodic orbits, SIAM J. Appl. Math. **58**, 227–234 (1998).

[5] K. Gopalsamy, Stability and Oscillations in Delay Differential Equations of Population Dynamics, Kluwer Academic Publishers, Dordrecht, 1992.

[6] K. Gopalsamy and R. A. Ahlip, Time delays in N-species competition—I: global stability in constant environments, Bull. Austral. Math. Soc. **27**, 427–441 (1983).

[7] Z. Hou, Global attractor in autonomous competitive Lotka-Volterra systems, Proc. Amer. Math. Soc. **127**, 3633–3642 (1999).

[8] Z. Hou, Global attractor in retarded competitive Lotka-Volterra systems, Nonlinear Analysis **47**, 4037–4048 (2001).

[9] Z. Hou, Global attractor in competitive Lotka-Volterra systems with retardation, Nonlinear Differ. Equ. Appl. **9**, 397–417 (2002).

[10] M. Kaykobad, Positive solutions of positive linear systems, Lin. Alg. Appl. **64**, 133–140 (1985).

[11] B. Li, S. Yu and X. Zeng, Some new results of autonomous Lotka-Volterra N-species competitive systems, Acta Math. Appl. Sin. **27**, 556–564 (2004).

[12] F. Montes de Oca and M. L. Zeeman, Balancing survival and extinction in nonautonomous competitive Lotka-Volterra systems, J. Math. Anal. Appl. **192**, 360–370 (1995).

[13] M. L. Zeeman, Hopf bifurcations in competitive three-dimensional Lotka-Volterra systems, Dynamics and Stability of Systems **8**, 189–217 (1993).

[14] M. L. Zeeman, Extinction in competitive Lotka-Volterra systems, Proc. Amer. Math. Soc. **123**, 87–96 (1995).

London Metropolitan University
Email: z.hou@londonmet.ac.uk